

References

1. Bucher PJ, Ijsselmuiden CB. Prevalence and causes of blindness in the northern Transvaal. *Br J Ophthalmol* 72: 721-6, 1988.
2. Chirambo MC, Schwab L. Ophthalmology in Malawi. *Arch Ophthalmol* 107: 1826-9, 1989.
3. Fafowora OF, Osuntokun OO. Age-related eye disease in the elderly members of rural African community. *East Afr Med J* 74: 435-7, 1997.
4. LeWallen S, Courtright P. Blindness in Africa: present situation and future needs. *Br J Ophthalmol* 85: 897-903, 2001.
5. Munoz B, West SK. Blindness and visual impairment in the Americas and the Caribbean. *Br J Ophthalmol* 86: 498-504, 2002.
6. Quigley HA. Number of people with glaucoma worldwide. *Br J Ophthalmol* 80: 389-393, 1996.
7. Silva JC, Bateman JB, Contreras F. Eye disease and care in Latin America and the Caribbean. *Surv Ophthalmol* 47: 267-74, 2002.
8. Thylefors B, Negrel AD. The global impact of glaucoma. *Bull World Health Organ* 72: 323-6, 1994.
9. Friedman DS. AGS Conference, San Francisco, 2003.
10. Sommer A, Tielsch JM, Katz J, et al. Racial differences in the cause-specific prevalence of blindness in east Baltimore. *N Engl J Med* 325:1412-7, 1991.
11. Tielsch JM, Sommer A, Katz J, et al. Racial variations in the prevalence of primary open-angle glaucoma. The Baltimore Eye Survey. *JAMA* 266: 369-74, 1991.
12. Martin MJ, Sommer A, Gold EB, Diamond EL. Race and primary open-angle glaucoma. *Am J Ophthalmol* 99: 383-7, 1985.
13. Wilson R, Richardson TM, Hertzmark E, Grant WM. Race as a risk factor for progressive glaucomatous damage. *Ann Ophthalmol* 17:653-9, 1985.
14. Weinreb RN. Ophthalmology August 2003.
15. Keltner JL, Johnson CA, et al. Confirmation of visual field abnormalities in the ocular hypertension treatment study. *Arch Ophthalmol*. 118: 1187-1194, 2000.
16. 1st Global AIGS Consensus Meeting on "Structure and Function in the Management of Glaucoma", San Diego, November 2003.
17. Kass MA, Heuer DK et al. The ocular hypertensive treatment study. A randomized trial determines that topical ocular hypotensive medication delays or prevents the onset of primary open-angle glaucoma. *Arch Ophthalmol*. 120: 701-713, 2002.
18. Collaborative Normal Tension Glaucoma Study Group. Natural history of normal tension glaucoma. *Ophthalmology*. 108: 247-253, 2001.
19. Liu JHK, Zhang X, Kripke DF, Weinreb RN. Twenty-four hour intraocular pressure pattern associated with early glaucomatous changes. *Invest Ophthalmol Vis Sci*. 44: 1586-1590, 2003.
20. Doughty MJ, Zaman ML. Human corneal thickness and its impact on intraocular pressure measures: a review and meta-analysis approach. *Surv of Ophthalmol*. 44: 367, 2000.
21. Leske MC, Connell AM, Wu SY, et al. Distribution of intraocular pressure. The Barbados Eye Study. *Arch Ophthalmol*. 115: 1051-7, 1997.
22. Quigley HA. Histopathology underlying glaucomatous damage II. 1st Global AIGS Consensus Meeting on "Structure and Function in the Management of Glaucoma", San Diego, November 2003.
23. Caprioli J., et al. Optic Disc Photographs. 1st Global AIGS Consensus Meeting on "Structure and Function in the Management of Glaucoma", San Diego, November 2003.
24. Quigley HA, Katz J, Derick RJ, Gilbert D, Sommer A. An evaluation of optic disc and nerve fiber layer examinations in monitoring progression of early glaucoma damage. *Ophthalmology* 99: 19-28, 1992.
25. Heijl A, Leske C, Bengtsson B, Hyman L, Bengtsson B, Hussein M. Reduction of intraocular pressure and glaucoma progression. *Arch Ophthalmol*. 120: 1268-1279, 2002.
26. Abrams LS, Scott IU, Spaeth GL, Quigley HA, Varma R. Agreement among optometrists, ophthalmologists, and residents in evaluating the optic disc for glaucoma. *Ophthalmology* 101: 1662-1667, 1994.
27. O'Conner DJ, Zeyen T, Caprioli, J. Comparisons of methods to detect glaucomatous optic nerve damage. *Ophthalmology* 100: 1498-1503, 1993.
28. Schultz RO, Radius RL, Hartz AJ, et al. Screening for glaucoma with stereo disc photography.

- Journal of Glaucoma* 4: 177-182, 1995.
29. Heijl A, Molder H. Optic disc diameter influences the ability to detect glaucomatous disc damage. *Acta Ophthalmol* 71: 122-129, 1993.
 30. Wollstein G, Garway-Heath DF, Fontana L, Hitchings RA. Identifying early glaucomatous changes. Comparison between expert clinical assessment of optic disc photographs and confocal scanning ophthalmoscopy. *Ophthalmology* 107: 2272-2277, 2000.
 31. Quigley HA, Addicks EM, Green WR. Optic nerve damage in human glaucoma. III. Quantitative correlation of nerve fiber loss and visual field defect in glaucoma, ischemic optic neuropathy, papilledema, and toxic neuropathy. *Arch Ophthalmol*. 100: 135-146, 1982.
 32. Quigley HA, Dunkelberger BS, Green WR. Retinal ganglion cell atrophy correlated with automated perimetry in human eyes with glaucoma. *AM J Ophthalmol*. 1989; 107: 453-464.
 33. Sommer A, Katz J, Quigley HA, et al. Clinically detectable nerve fiber atrophy precedes the onset of glaucomatous field loss. *Arch Ophthalmol*. 1991; 109: 77-83.
 34. Sommer A, Miller NR, Pollack I, et al. The nerve fiber layer in the diagnosis of glaucoma. *Arch Ophthalmol*. 1977; 95: 2149-56.
 35. Tuulonen A, Lehtola J, Airaksinen PJ. Nerve fiber layer defects with normal visual fields. Do normal optic disc and normal visual field indicate absence of glaucomatous abnormality? *Ophthalmology* 1993; 100: 587-598.
 36. Airaksinen PJ, Mustonen E, Alanko HI. Optic disc hemorrhages precede retinal nerve fiber layer defects in ocular hypertension. *Acta Ophthalmol*. 59: 627-41, 1981.
 37. Caprioli J, Prum B, Zeyen T. Comparison of methods to evaluate the optic nerve head and nerve fiber layer for glaucomatous damage. *Am J Ophthalmol*. 121: 659-67, 1996.
 38. Sample PA, Bosworth CF, Blumenthal EZ, Girkin C, Weinreb RN. Visual function-specific perimetry for indirect comparison of different ganglion cell populations in glaucoma.
 39. Wang F, Tielsch JM, Ford DE, Quigley HA, Whelton PK. Evaluation of screening schemes for eye disease in a primary care setting. *Ophthalmic Epidemiol* 5: 69-82, 1998.
 40. Hoyt WF, Newman NM. The earliest observable defect in glaucoma? *Lancet*. 1972; 1: 692-693.
 41. Airaksinen PJ, Alanko HI. Effect of retinal nerve fiber loss on the optic nerve head configuration in early glaucoma. *Graefes Arch Clin Exp Ophthalmol* 220: 193-196, 1983.
 42. Quigley HA. Examination of the retinal nerve fiber layer in the recognition of early glaucomatous damage. *Trans Am Ophthalmol Soc*. 84: 920-66, 1986.
 43. Quigley HA, Enger C, Katz J, Sommer A et al. Risk factors for the development of glaucomatous visual field loss in ocular hypertension. *Arch Ophthalmol* 112: 644-9, 1994.
 44. Jonas JB, Hayreh SS. Localised retinal nerve fiber layer defects in chronic experimental high pressure glaucoma in rhesus monkeys. *Br J Ophthalmol* 83: 1291-5, 1999.
 45. Tuulonen A, Airaksinen PJ. Initial glaucomatous optic disc and retinal nerve fiber layer abnormalities and their progression. *Am J Ophthalmol* 1991; 111: 485-490.
 46. Quigley HA, Miller NR, George T. Clinical evaluation of nerve fiber layer atrophy as an indicator of glaucomatous optic nerve damage. *Arch Ophthalmol*. 98: 1564-71, 1980.
 47. Wang F, Quigley HA, Tielsch JM. Screening for glaucoma in a medical clinic with photographs of the nerve fiber layer. *Arch Ophthalmol*. 112: 796-800, 1994.
 48. Paczka JA, Quigley HA, Friedman DS, Barron Y, Vitale S. Diagnostic capabilities of frequency-doubling technology, scanning laser polarimetry and nerve fiber layer photographs to distinguish glaucomatous damage. *Am J Ophthalmol*. 131: 188-197, 2001.
 49. Airaksinen PJ, Drance SM, et al. Diffuse and localized nerve fiber loss in glaucoma. *Am J Ophthalmol*. 98: 566-71, 1984.
 50. Zhou Q, Weinreb RN. Individualized compensation of anterior segment birefringence during scanning laser Polarimetry. *Invest Ophthalmol Vis Sci*. 2002; 43: 2221-2228.
 51. Blumenthal EZ, Frenkel S. Inter-device reproducibility of the scanning laser polarimeter with variable cornea compensation. *Eye* in press, 2004.
 52. Shirakashi M, Yaoeda K, Fukushima A, Funaki S, Funaki H, Ofuchi N, Abe H. Retinal nerve fiber layer thickness measurement with the new GDx Access. *Journal of the Eye* 20: 407-409, 2003.
 53. Reus NJ, Lemij HG. Sensitivity and specificity of the GDx VCC. *International Society for Imaging in the Eye* conference. Fort Lauderdale, FL. 2003.
 54. Medeiros F, Zangwill LM, Bowd C, Mohammadi, K, Weinreb RN. Comparison of scanning laser po-

- larimetry using variable corneal polarization compensation and retinal nerve fiber layer photography for detection of glaucoma. *Arch Ophthalmol*. In press, 2004.
55. Weinreb RN, Dreher AW, Coleman A, Quigley HA, Shaw B, Reiter K. Histopathologic validation of Fourier-ellipsometry measurements of retinal nerve fiber layer thickness. *Arch Ophthalmol*. 108: 557-560, 1990.
 56. Morgan JE, Waldock A, Jeffery G, Cowey A. Retinal nerve fiber layer polarimetry: histological and clinical comparison. *Br J Ophthalmol*. 82: 684-690, 1998.
 57. Dreher AW, Reiter K. Scanning laser polarimetry of the retinal nerve fiber layer. *Proc SPIE* 1746: 34-41, 1992.
 58. Dreher AW, Reiter K. Retinal laser ellipsometry: a new method for measuring the retinal nerve fiber thickness distribution? *Clin Vision Sci*. 7: 481-488, 1992.
 59. Dreher AW, Reiter K, Weinreb RN. Spatially resolved birefringence of the retinal nerve fiber layer assessed with a retinal laser ellipsometer. *Appl Opt*. 31: 3730-3735, 1992.
 60. Zhou Q, Reed J, et al. Detection of glaucomatous retinal nerve fiber layer damage by scanning laser polarimetry with variable corneal compensation. *Proc SPIE* 4951: 1-10, 2003.
 61. Weinreb RN, Bowd C, Greenfield DS, Zangwill LM. Measurement of the magnitude and axis of corneal polarization with scanning laser polarimetry. *Arch Ophthalmol*. 2002; 120: 901-906.
 62. Greenfield DS, Knighton RW, Feuer W, Schiffman J, Zangwill L, Weinreb RN. Correction for corneal polarization axis improves the discriminating power of scanning laser polarimetry. *Ophthalmology* 108: 1065-1069, 2002.
 63. Garway-Heath DF, Greaney MJ, Caprioli J. Correction for the erroneous compensation of anterior segment birefringence with the scanning laser polarimeter for glaucoma diagnosis. *Invest Ophthalmol Vis Sci*. 43: 1465-1474, 2002.
 64. Knighton RW, Huang X-R. Analytical methods for scanning laser polarimetry. *Optics Express* 20: 1179-1189, 2002.
 65. Bagga H, Greenfield DS, Knighton RW. Scanning laser polarimetry with variable corneal compensation: identification and correction for corneal birefringence in eyes with macular disease. *Invest Ophthalmol Vis Sci*. 44: 1969-1976, 2003.
 66. Morrison JC, Cork LC, Dunkelberger GR, Brown A, Quigley HA. Aging changes of the rhesus monkey optic nerve. *Invest Ophthalmol Vis Sci*. 31: 1623-1627, 1990.
 67. Balazsi AG, Rootman J, Drance SM, et al. The effect of age on the nerve fiber population of the human optic nerve. *Am J Ophthalmol* 97: 760-766, 1984.
 68. Gao H, Hollyfield JG. Aging of the human retina. *Invest Ophthalmol Vis Sci*. 33: 1-17, 1992.
 69. Bowd C, Zangwill LM, Blumenthal EZ, Vasile C, et al. Imaging of the optic disc and retinal nerve fiber layer: the effect of age, optic disc area, refractive error, and gender. *J Opt Soc Am A Opt Image Sci Vis*. 19: 197-207, 2002.
 70. Toprak AB, Yilmaz OF. Relation of optic disc topography and age to thickness of retinal nerve fiber layer as measured using scanning laser polarimetry, normal subjects. *Br J Ophthalmol*. 84: 473-8, 2000.
 71. Weinreb RN, Shakiba S, Sample PA, et al. Association between quantitative nerve fiber layer measurement and visual field loss in glaucoma. *Am J Ophthalmol* 120: 732-738, 1995.
 72. Chi Q, Tomita G, Inazumi K, Hayakawa T, Tadayoshi I, Kitazawa Y. Evaluation of the effect of aging on the retinal nerve fiber layer thickness using scanning laser polarimetry. *J Glaucoma* 4: 406-413, 1995.
 73. Poinosawmy D, Fontana L, Wu JX, Fitzke FW, Hitchings RA. Variation of nerve fiber thickness measurements with age and ethnicity by scanning laser polarimetry. *Br J Ophthalmol*. 81: 350-354, 1997.
 74. Reus NJ, Colen TP, Lemij HG. Visualization of localized retinal nerve fiber layer defects with the GDx with individualized and with fixed compensation of anterior segment birefringence. *Ophthalmology* 110: 1512-1516, 2003.
 75. Weinreb RN, Bowd C, Zangwill LM. Scanning laser polarimetry in monkey eyes using variable corneal polarization compensation. *Journal of Glaucoma* 11: 378-84, 2002.
 76. Weinreb RN, Bowd C, Zangwill LM. Glaucoma detection using scanning laser polarimetry with variable corneal compensation. *Arch Ophthalmol* 121: 218-224, 2003.

77. Medeiros FA, Zangwill LM, Bowd C, Bernd AS, Weinreb RN. Fourier analysis of scanning laser polarimetry measurements with variable corneal compensation in glaucoma. *Invest Ophthalmol Vis Sci.* 44: 2606-2612, 2003.
78. Shirakashi M, Yaoeda K, Fukushima A, Funaki S, Funaki H, Ofuchi N, Nakatsue T, Abe H. Usefulness of GDx VCC in glaucoma detection. *Journal of the Eye* 20: 1019-1021, 2003.
79. Bowd C, Zangwill LM, Weinreb RN. Association between scanning laser polarimetry measurements using variable corneal polarization compensation and visual field sensitivity in glaucomatous eyes. *Arch Ophthalmol* 121: 961-966, 2003.
80. Bagga H, Greenfield DS, Feuer W, Knighton RW. Scanning laser polarimetry with variable corneal compensation and optical coherence tomography in normal and glaucomatous eyes. *Am J Ophthalmol* 135: 521-529, 2003.
81. Garway-Heath DF, Greenfield DS, Caprioli J. Correlation of visual field sensitivity and retinal nerve fiber layer thickness as measured by scanning laser polarimetry. *Invest Ophthalmol Vis Sci, Suppl.* 980, 2003.
82. Tjon-Fo-Sang MJ, Lemij HG. The sensitivity and specificity of nerve fiber layer measurements in glaucoma as determined with scanning laser polarimetry. *Am J Ophthalmol.* 123: 62-69, 1997.
83. Poinoosawmy D, Tan JC, Bunce C, Hitchings RA. The ability of the GDx nerve fibre analyser neural network to diagnose glaucoma. *Graefes Arch Clin Exp Ophthalmol.* 239: 122-7, 2001.
84. Sinai MJ, Essock EA, Fechtner RD, Srinivasan N. Diffuse and localized nerve fiber layer loss measured with a scanning laser polarimeter: sensitivity and specificity of detecting glaucoma. *J Glaucoma* 9: 154-162, 2000.
85. Essock EA, Sinai MJ, Bowd C, Zangwill LM, Weinreb RN. Fourier analysis of optical coherence tomography and scanning laser polarimetry retinal nerve fiber layer measurements in the diagnosis of glaucoma. *Arch. Ophthalmol.* 121: 1238-45, 2003.
86. Colen TP, Lemij HG. Sensitivity and specificity of the GDx: clinical judgment of standard printout versus the number. *J Glaucoma* 12: 129-33, 2003.
87. Zangwill LM, Bowd C, Berry CC, Williams J, Blumenthal EZ, Sanchez-Galeana CA, Vasile C, Weinreb RN. Discriminating between normal and glaucomatous eyes using the Heidelberg Retina Tomograph, GDx Nerve Fiber Analyzer, and Optical Coherence Tomograph. *Arch Ophthalmol.* 119: 985-993, 2001.
88. Tribble JR, Schultz RO, Robinson JC, Rothe TL. Accuracy of scanning laser polarimetry in the diagnosis of glaucoma. *Arch Ophthalmol.* 117: 1298-1304, 1999.
89. Choplin NT, Lundy DC. The sensitivity and specificity of scanning laser polarimetry in the detection of glaucoma in a clinical setting. *Ophthalmology* 108: 899-904, 2001.
90. Bowd C, Zangwill LM, Berry CC, Blumenthal EZ, Vasile C, Sanchez-Galeana C, Bosworth CF, Sample PA, Weinreb RN. Detecting early glaucoma by assessment of retinal nerve fiber layer thickness and visual function. *Invest Ophthalmol Vis Sci.* 42: 1993-2003, 2001.
91. Weinreb RN, Zangwill LM, Berry CC, Bathija R, Sample PA. Detection of glaucoma with scanning laser polarimetry. *Arch Ophthalmol.* 116: 1583-589, 1998.